

Terrazole®

35% WETTABLE POWDER

TURF & ORNAMENTAL FUNGICIDE

Active Ingredients: (% by weight)
Etridiazole [5-Ethoxy-3-(trichloromethyl)-1,2,4-thiadiazole]35%
Other Ingredients:65%
Total: 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF IN EYES:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor.• Do not give anything by mouth to an unconscious person.
IF ON SKIN OR ON CLOTHING:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15 to 20 minutes.• Call a poison control center or doctor for treatment advice.
EMERGENCY ASSISTANCE: Have the product container or label with you when calling a poison control center or doctor, or going for treatment.	
24 Hour Emergency Number 1-866-928-0789	
Product Use Information Number: 1-866-761-9397	

NET CONTENTS:

Manufactured for:
MacDermid Agricultural Solutions, Inc.
245 Freight Street
Waterbury, CT 06702-1818

EPA REG. NO. 70506-569-59807
EPA EST. NO.
019/062608

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC
ANIMALS CAUTION

Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical resistance category selection chart.

All mixers, loaders, applicators and other handlers must wear: coveralls over long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material, chemical resistant footwear plus socks, NIOSH approved respirator with: an organic-vapor removing cartridge with a prefilter approved for pesticides (MSHA/NIOSH approval number prefix TC-23C), or a canister approved for pesticides (MSHA/NIOSH approval number prefix TC-14G), or a NIOSH approved respirator with an OV cartridge or canister with any N2, R, P or HE prefilter, chemical-resistant apron when mixing, loading, cleaning equipment.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables, use detergent and hot water.

Keep and wash PPE separately from other laundry.

When all other handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR 170.240(d)(5)), the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Surface Water Advisory: Etridiazole can contaminate surface water through spray drift. Under some conditions, etridiazole may have a high potential for runoff into surface water for several weeks post application. These conditions include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with infield canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain surface water.

PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product to turf through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not allow this product to drift. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

INDOOR RESTRICTIONS: Entry (including early entry that would otherwise be permitted under the WPS) into greenhouses, potting sheds, and other indoor areas by any person - other than a correctly trained applicator who is performing a handling task permitted by the WPS and who is wearing the required handler PPE including a respirator - is **PROHIBITED** in the entire enclosed structure/building from the start of application until application is complete and one of the following ventilation criteria (providing outside air) is met: (1) 10 air exchanges; (2) 2 hours of fans or other mechanical ventilation providing outside air; (3) 4 hours of vents, windows, or other passive ventilation; (4) 11 hours with no ventilation followed by 1 hour of mechanical ventilation; (5) 11 hours of no ventilation followed by 2 hours of passive ventilation; or (6) 24 hours with no ventilation. After ventilation criteria are met and until the REI expires, do not enter or allow worker entry into treated areas, except as provided in the WPS. Note: after the expiration of the REI whenever Terrazole- treated soil or planting media is being handled or disturbed indoors, continuous ventilation of the area is required at a minimum rate of one complete air exchange per hour.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- long-sleeved shirt and long pants
- chemical-resistant gloves made of any waterproof material
- shoes and socks.
- protective eyewear

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40CFR part 170). WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Re-entry Statement:

Do not enter or allow others to enter until sprays have dried.

GENERAL INFORMATION

TERRAZOLE 35WP soil fungicide is effective for prevention and control of damping-off, root rot and stem diseases caused by *Pythium* and *Phytophthora* species in golf course turf, ornamentals and nursery crops.

MIXING INSTRUCTIONS

Fill the mixing tank with half the required amount of water. Then add TERRAZOLE 35WP with agitation running to insure proper mixing and suspension of product. Fill the tank with the remaining amount of required water.

The mixture of TERRAZOLE 35WP and water should be continuously agitated to assure uniform application.

COMPATIBILITY: For broad spectrum disease control, this product can be tank mixed with TERRACLO® 75% WP (EPA Reg. No. 400-399) or TERRACLO® 400 (EPA Reg. No. 400-454) for use on ornamentals in accordance with the more restrictive label limitations and precautions. No label dosage rates may be exceeded. This product cannot be mixed with any product containing a label prohibition against such mixing. Do not mix with other chemicals unless prior use has proven compatibility.

GOLF COURSE TURF - TEES AND GREENS

TERRAZOLE 35WP is effective for the control of *Pythium* blight, cottony blight, grease spot, spot blight, root rot, crown rot and damping-off disease caused by *Pythium* spp.

TERRAZOLE 35WP can be applied as a spray to both newly seeded and established turf.

Application is limited to tees and greens. Application to fairways is prohibited.

For commercial use only. Not for use on home lawns, sod farms or municipal parks.

Do not allow animals to graze on treated areas and do not feed clippings from treated areas to animals.

USE AND RATE RECOMMENDATIONS

Mix 2 to 4 ozs. in 2 to 5 gallons of water and apply to 1000 sq. ft. Begin applications at first sign of disease and retreat at 10 to 14 day intervals as needed.

Within the rate range given, use the lower rate for the shortest interval listed and the higher rate for the longest interval.

Under severe conditions, use the highest rate and shortest interval specified.

Do not exceed 4.0 ozs. per 1000 sq. ft. per application or 10 ozs. per 1000 sq. ft. per year.

The interval between applications may not be less than 10 days.

ORNAMENTALS

TERRAZOLE 35WP controls water mold, damping off, root rot, stem rot and crown rot diseases caused by *Pythium* spp. and *Phytophthora* spp. TERRAZOLE 35WP is mixed with water and applied as a soil drench, beginning at the time of seeding or transplanting. For use in commercial nurseries and greenhouses only. Use in residential greenhouses or other indoor plant sites is prohibited.

APPLICATION NOTES

Apply the recommended rate of TERRAZOLE 35WP according to the guidelines below. The drench solution should be continually agitated to assure uniform application of the fungicide material. To improve penetration of the fungicide, the drenching must be followed with an additional quantity of water equal to at least half the volume of the fungicidal drench, applied either by sprinkling or irrigation.

For Bed and Bench Grown Plants:

Soil Depth (inches)	Sq.ft./100 gals.	Fl.oz./Sq.ft.
3 or less	800	16
4	600	21
5	480	27
6	400	32
7	340	37
8	300	43
9	270	48
10	240	53
12 or more	200	64

For Container Grown Plants:

Pot Diameter (inches)	Minimum Drench Volume fl.oz./pot
4	2
5	3
6	4
8	10
10	20
12	30

USE AND RATE RECOMMENDATIONS

CROP	OUNCES TERRAZOLE 35WP PER 100 GALS. (TSP./GAL)	REMARKS
Bedding plants Flowering plants Foliage plants Woody ornamentals	3.5 - 10 (½ - 1 ½)	Re-treat at 4 to 12 week intervals as necessary. Use higher rates for peat and other high organic potting media.

PLANT TOLERANCE:

Neither the manufacturer nor the seller has determined whether or not TERRAZOLE 35WP can be used safely on all ornamental plants. Prior to any large scale application on such plants, the user should determine the safety of TERRAZOLE 35WP fungicide by testing a small number of the type of plants to be treated at the recommended rates for that particular group, i.e., bedding plants, foliage, etc.

USE DIRECTIONS FOR CHEMIGATION:

In addition to the above use rates and recommendations, the following precautions must be observed when using this product in any type of irrigation system:

Apply this product only through the following systems:

1) Pressurized drench (flood) of drip (trickle) systems, 2) Micro-irrigation such as spaghetti-tube or individual tube irrigation, 3) Hand-held calibrated irrigation equipment such as the hand-held wand with injector, 4) Ebb and flow systems.

Do not apply this product through any other type of irrigation system.

The Terrazole mixture should be continually agitated to assure uniform application of the fungicide material. To improve penetration of the fungicide, the drenching must be followed with an additional quantity of water equal to at least half the volume of the fungicidal drench, applied either by sprinkling or irrigation.

Crop injury or lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water.

If you have any questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts. Do not connect an irrigation system, (including greenhouse systems), used for pesticide application to a public water system unless the pesticide label prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

SPRINKLER CHEMIGATION

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

SYSTEMS CONNECTED TO PUBLIC WATER SYSTEMS

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days of the year.

Chemigation systems connected to public water systems must contain a functional, reduced pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line up-stream from the point of pesticide introduction. As an option to the RPZ, the water from the public water systems should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where the pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store in a dry, secure location.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Paper Bags: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Completely empty into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT NOTICE — Seller warrants that this product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with the directions and instructions specified on the label under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product, contrary to label instructions, or under conditions not reasonably foreseeable to seller and, to the extent consistent with applicable law, the buyer assumes the risk of any such use.

®TERRAZOLE is a Registered Trademark of MacDermid Agricultural Solutions, Inc.

®TERRACLOR is a Registered Trademark of AMVAC Corporation

©Copyright 2015, MacDermid Agricultural Solutions, Inc.