

SHUTTLE™ O

Miticide



FOR USE ON GREENHOUSE, SHADEHOUSE, AND FIELD-GROWN ORNAMENTAL, FLORAL, FOLIAGE, AND NURSERY CROPS, AND INTERIORSCAPES

Product Information Bulletin

7 2 5

GENERAL INFORMATION

SHUTTLE™ O miticide is the first product from the naphthoquinone derivative class of chemistry (MOA Group 20B). Because of its unique chemical class, **SHUTTLE O** is an excellent choice as a rotational tool for control of two-spotted spider mites, spruce spider mites, and other mites (see box at bottom).

SHUTTLE O is a suspension concentrate formulation containing 1.25 pounds of active ingredient acequinocyl per gallon.

Common Name: Acequinocyl

Class: Naphthoquinone derivative

GROUP	20B	INSECTICIDE
-------	-----	-------------

Packaging: 16 fluid ounce (pint) container, 4 per case

RESTRICTED ENTRY INTERVAL (REI) AND SIGNAL WORD

REI = **12 hours**

Signal Word = **CAUTION**

PERSONAL PROTECTIVE EQUIPMENT (PPE)

- long-sleeved shirt and long pants
- socks
- shoes
- chemical resistant gloves made of water proof material

ADVANTAGES OF SHUTTLE O

- Long-term residual control (21 days)
- Unique mode of action with no cross-resistance
- Control of all mite life stages (egg, larvae, nymphs, adults)
- Virtually harmless to predacious mites and beneficial insects
- Reduced Risk pesticide classification
- Broad use area labeling – inside, outside, interiorscapes

MODE OF ACTION

SHUTTLE O is effective on all life cycle stages of two-spotted spider mites and spruce spider mites through both contact and ingestion. **SHUTTLE O** works by interfering with energy production in the mitochondria of the target pest's cells.

SHUTTLE O features a unique mode of action with no cross-resistance to other miticides. It is an ideal fit in a mite resistance management strategy.

USE SITES

SHUTTLE O may be used on ornamentals grown in greenhouses, container and field nurseries, lath and shade houses, arboretums, and interiorscapes.

TARGET PESTS CONTROLLED

- Two spotted spider mite
- Spruce spider mites
- Citrus red mite
- European red mite
- Pacific spider mite
- Willamette spider mite
- Strawberry spider mite
- Texas citrus mite

APPLICATION AND RATES

SHUTTLE O application rates per 100 gallons of water: 6.4 to 12.8 fluid ounces (.06 to .125 pounds of active ingredient/100 gallons). **SHUTTLE O** should be applied when threshold mite populations are observed and before large mite populations have been established.

Apply **SHUTTLE O** in sufficient water to ensure uniform and thorough coverage of foliage.

Users should not apply less than 100 gallons of **SHUTTLE O** finished spray per acre.

TANK MIXING

SHUTTLE O can be combined with most commonly-used pesticides with the exception of strongly alkaline materials, which may reduce the activity of the product. Do not combine **SHUTTLE O** with Aliette® WDG (fosetyl).

For broad-spectrum control, **SHUTTLE O** may be tank mixed with most other miticide/insecticide products.

SHUTTLE O and various tank mix combinations including spreader stickers and adjuvants, have not been tested on all plant species and the many varieties of each.

Tank mix combinations should be trialed on a small-scale basis before any large-scale use.

RESISTANCE MANAGEMENT

When used as directed, **SHUTTLE O** is effective on a wide variety of mite species. **SHUTTLE O** shows efficacy on all mite life cycle stages.

Users should not make successive applications of **SHUTTLE O**. Users should apply **SHUTTLE O** as

part of a resistance management strategy program that includes rotation with other miticides with different modes of action.


RESTRICTIONS & LIMITATIONS

- **SHUTTLE O** should not be used through any type of irrigation system.
- **SHUTTLE O** should not be applied by air.
- Do not apply **SHUTTLE O** within 75 feet of aquatic areas.
- Do not apply to impatiens (including New Guineas) or miniature roses without first testing for sensitivity.


SHUTTLE O provides excellent control in university and laboratory testing

Even when scouted, mite populations can explode quickly and damage ornamental plants. That's why the activity of **SHUTTLE O** on all life cycle stages is helpful to growers.

- **SHUTTLE O** delivers powerful contact activity for control of all mite life stages.
- Continues to work through ingestion to keep mite populations in check for three weeks or more.



Spray Volume: 100 GPA; Application Date: 8/24/2005; Evaluation: 8/31, 9/7, 9/28; DAT = Days after treatment



Spray Volume: 100 GPA; Application Date: 6/1/2002; Evaluation: 6/29, Control (No./4 Leaves): Adults, 60; Nymphs, 65; Eggs, 136