

## FIFRA Section 2(ee) Recommendation

**For Control of Rice Root Aphid and Root Mealybug on Listed Food and Nonfood Crops Grown in Indoor/Outdoor Nurseries, Greenhouses, Shadehouses, and Interiorscapes**

**FOR USE AND DISTRIBUTION IN THE STATE OF TEXAS ONLY**

**Expires 12/31/2026**

***This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by the US Environmental Protection Agency.***

All applicable directions, restrictions, and precautions on the EPA-registered labels must be followed. Please contact your Certis USA regional sales manager to determine the specific requirements for FIFRA §2(ee) recommendations in your respective state.

### **DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

#### ***All Listed Crops***

Insect	Application Rate	Use Instructions
Rice Root Aphid	Spray: 1-2 lb per 100 gallons spray volume	High pest pressure: apply 2-3 times, 3-5 days apart Low pest pressure: apply every 7-10 days  No limit to application number in a single season
	Dip: ¼ - ½ oz per gallon (2.5 – 5.0 oz per 10 gallons)	Agitate dip suspension throughout use. Prepare a new dip suspension regularly. Prepare only as much dip suspension as can be used in one day. Do not use dip suspension for more than one day. Disinfect the dipping tank and equipment before a new dip suspension is prepared.  Avoid exposing dipped cuttings to full sun. Do not use on stressed/wilting cuttings. Do not use without determining crop sensitivity (phytotoxicity) first.



<p>Rice Root Aphid</p> <p>Root mealybug</p>	<p>Drench:  <math>\frac{1}{4}</math> - <math>\frac{1}{2}</math> oz per gallon          (2.5 – 5.0 oz per 10          gallons)</p>	<p>Agitate drench suspension throughout use. Prepare only as much drench suspension as can be used in one day. Do not use drench suspension for more than one day.</p> <p>Ensure growing medium is moist prior to application to facilitate the movement of spores throughout the substrate. Apply sufficient volume to ensure thorough wetting of the growing medium but not to the point of leaching. Ensure uniform distribution of applied product.</p>
<p><b>Crop-Specific Recommendation:</b> Hemp foliar applications should be suspended 3-4 weeks prior to harvest.</p>		

**Crop-Specific Restrictions:**

Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

Do not use dip solution for more than one day.

Do not apply through a thermal pulse fogger.

Refer to EPA-approved label for complete list of **USE RESTRICTIONS**.

This FIFRA Section 2(ee) recommendation contains new or additional directions for use of this product, which may not appear on the package label. Read and observe the precautionary statements plus all other information appearing on the product labels. **This recommendation must be in the possession of the user at the time of pesticide application.**

